

Bibliografia

- Abrams P, Cardozo L, Fall M, et al. *The standardisation of terminology of lower urinary tract function: report from the Standardisation Sub-committee of the International Continence Society*. *Neurourol Urodyn* 2002;21:167-78.
- Abrams P, Cardozo L, Wagg A, Wein, A. (Eds) *Incontinence 6th Edition* (2017). ICI-ICS. International Continence Society, Bristol UK, ISBN: 978-0956960733.
- Callegari E, Malhotra B, Bungay PJ, et al. A comprehensive non-clinical evaluation of the CNS penetration potential of antimuscarinic agents for the treatment of overactive bladder. *Br J Clin Pharmacol*. 2011;72(2):235-46.
- Chapple CR, Khullar V, Gabriel Z, Muston D, Bitoun CE, Weinstein D. The effects of antimuscarinic treatments in overactive bladder: an update of a systematic review and meta-analysis. *Eur Urol*. 2008 Sep;54(3):543-62.
- Kurpad R, Kennelly MJ. The evaluation and management of refractory neurogenic overactive bladder. *Curr Urol Rep*. 2014;15(10):444.
- Lucas MG, Bedretidnova D, Bosch JLHR, et al. *Guidelines on Urinary Incontinence*. © European Association of Urology 2014. Disponibile su https://uroweb.org/wp-content/uploads/20-Urinary-Incontinence_LR.pdf (ultimo accesso: 28/02/2019)
- Madhuvrata P, Cody JD, Ellis G, Herbison GP, Hay-Smith EJ. Which anticholinergic drug for overactive bladder symptoms in adults. *Cochrane Database Syst Rev*. 2012 Jan 18;1:CD005429.
- McDonagh, et al., *Drug class review: agents for overactive bladder*. Final report Update 4. Portland (OR): Oregon Health & Science University; 2009 Drug Class Reviews.
- Nambiar A, Lucas M. Chapter 4: Guidelines for the diagnosis and treatment of overactive bladder (OAB) and neurogenic detrusor overactivity (NDO). *Neurourol Urodyn*. 2014;33 Suppl 3:S21-5.
- Risacher SL, McDonald BC, Tallman EF, et al. for the Alzheimer's Disease Neuroimaging Initiative. Association Between Anticholinergic Medication Use and Cognition, Brain Metabolism, and Brain Atrophy in Cognitively Normal Older Adults. *JAMA Neurol*. 2016;73(6):721-32.
- Sakakibara R, Panicker J, Finazzi-Agro E, et al; Parkinson's Disease Subcommittee, The Neurourology Promotion Committee in The International Continence Society. A guideline for the management of bladder dysfunction in Parkinson's disease and other gait disorders. *Neurourol Urodyn*. 2015; 35(5):551-63.
- Shamliyan TA, Kane RL, Wyman J, et al. Systematic review: randomized, controlled trials of nonsurgical treatments for urinary incontinence in women. *Ann Intern Med* 2008; 148(6): p. 459-73.
- Thüroff JW, Abrams P, Andersson KE, et al. EAU guidelines on urinary incontinence. *Eur Urol*. 2011;59(3):387-400.
- Vouri SM, Schootman M, Strobe SA, Xian H, Olsen MA. Antimuscarinic use and discontinuation in an older adult population. *Arch Gerontol Geriatr*. 2019 Jan - Feb;80:1-11. doi: 10.1016/j.archger.2018.09.005. Epub 2018 Sep 22. PubMed PMID: 30268971.
- Vouri SM, Kebodeaux CD, Stranges PM, Teshome BF. Adverse events and treatment discontinuations of antimuscarinics for the treatment of overactive bladder in older adults: A systematic review and meta-analysis. *Arch Gerontol Geriatr*. 2017;69:77-96.
- Yamada S, Ito Y, Nishijima S, Kadekawa K, Sugaya K. Basic and clinical aspects of antimuscarinic agents used to treat overactive bladder. *Pharmacol Ther*. 2018;189:130-148.

